
The Origin of E+A Galaxies

Tomotsugu GOTO and the SDSS Collaboration

Department of Astronomy, School of Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

1. E+A Mystery

E+A galaxies are galaxies with strong Balmer absorption lines and with no significant emission in [OII] or $H\alpha$ lines. The existence of strong Balmer absorption lines shows that these galaxies have experienced a starburst within the last 1 Gyr. At the same time, these galaxies do not show any sign of on-going star formation as indicated by the lack of [OII] or $H\alpha$ emission lines. Therefore, E+A galaxies are interpreted as a post-starburst galaxy — a galaxy which truncated starburst suddenly. However, for more than 20 years, the origin of this post-starburst phenomena remained a mystery. Using the 100,000 galaxy spectra of the Sloan Digital Sky Survey, we tested three popular explanations to the E+A phenomenon.

2. Are E+As Created by a Cluster-Related Mechanism?

At first, E+A galaxies were found in cluster regions. Therefore, cluster related physical mechanisms have been thought to be responsible for the E+A phenomena. However, in the left panel of Fig. 1, E+A galaxies show the same local galaxy density distribution as field (all) galaxies. These numerous field E+A galaxies cannot be explained with a cluster related physical mechanism.

3. Are E+As Dusty Starbursts?

Another possible explanation for E+A phenomena is a dusty starburst, where E+A galaxies are actually star-forming, but emission lines are invisible in optical wavelengths due to the heavy obscuration by dust. However, in the middle panel of Fig.1, all E+As have $r - K < 1.1$, where the theoretical models predict $r - K \sim 1.5$ for dusty starburst galaxies. In the left panel of Fig.1, no E+As show any sign of starburst in radio estimated (dust-free) star formation rate derived from the FIRST data.




Fig. 1. (left) Local galaxy density distribution for E+As and all galaxies. (middle) SDSS-2MASS optical-near infrared color distribution. (right) Radio estimated star formation rate derived from the FIRST data.

4. Are E+As Merger/Interaction Origin?

Merger/Interaction between two galaxies also can induce star formation. In Table 1, we counted number of accompanying galaxies within 50 and 75 kpc around E+A galaxies and compared those with randomly selected normal galaxies. We use galaxies with $-23 < Mr < -19.5$ and statistically subtract background counts using the global galaxy number counts in the SDSS data. As shown in the table, E+As have more accompanying galaxies than normal galaxies. Especially E+As with the strongest $H\delta$ absorption ($H\delta$ EW $> 7\text{\AA}$) have 8 times more accompanying galaxies within 50 kpc. Since this excess in the number of accompanying galaxies is significant, we conclude that E+As are most likely to have a merger/interaction origin [1][2].

Table 1. Number of accompanying galaxies around all the E+A galaxies and E+As with $H\delta$ EW > 7 are compared with that of 1000 randomly picked galaxies. We use galaxies with $-23.0 < Mr^* < -19.5$ after k -correction within the radius of 50 and 75 kpc. Fore/background galaxy number counts are statistically subtracted.

Sample	$N_{\text{nearby galaxy}}$ within 50 kpc	$N_{\text{nearby galaxy}}$ within 75 kpc
Random	0.03 ± 0.01	0.16 ± 0.01
All E+As	0.12 ± 0.03	0.26 ± 0.04
E+A with $H\delta$ EW $> 7\text{\AA}$	0.24 ± 0.09	0.40 ± 0.12

References

- [1] T. Goto, PhD Thesis, 2003, The University of Tokyo, astro-ph/0310196
- [2] T. Goto et al., 2003, PASJ, 55, 771